

Title of Invited Lecture

The author(s)'s name(s)*

Abstract. This is to explain how to prepare your Invited Lecture for publication in the Proceedings of the ICIAM '07. The abstract should be comprehensible to any mathematician. In general terms should be described what the paper is about. A rough statement in words is preferable to a precise statement loaded with symbols and technical notions.

Mathematics Subject Classification (2000). Primary 00A05; Secondary 00B10.

Keywords. General mathematics, collection of articles.

1. Introduction

Authors are requested to use standard L^AT_EX and the class file

`icmart.cls`

This file is essentially 'article.cls', slightly changed, loading `amsmath`, `amsfonts`, `amssymb`, `latexsym`, and with `amsthm.sty` included. It sets the page size to

```
\textheight=548pt
\textwidth=357pt
```

The T_EX source file should begin with

```
\documentclass{icmart}
```

No personal style files should be used. Each paper should contain the 2000 Mathematics Subject Classification. We would very much appreciate it if you could avoid one-letter lower case newly defined commands like

```
\def\epsilon{\varepsilon}
```

since this can interfere with conversion of your article to Times fonts later.

2. Some rules

In order to achieve a uniform appearance of all the contributions, we encourage you to observe the following rules when preparing your article.

*The authors are grateful to the Max Planck Institute (Bonn) for hospitality during the writing of this paper.

2.1. Displayed formulas. If you have displayed formulas consisting of more than one line we would prefer if you use

`\begin{align}...\end{align}`

instead of

`\begin{eqnarray}...\end{eqnarray}`

(respectively the starred forms) since the former yields a better spacing. Compare:

$$A = f(x_i) = F'(x) \tag{1}$$

$$B = g(x_i) = G'(x) \tag{2}$$

$$A = f(x_i) = F'(x) \tag{3}$$

$$B = g(x_i) = G'(x) \tag{4}$$

In case you do not want the numbering for every line, type

`\nonumber`

at the end of the line where you do not want a number.

$$\begin{aligned} A &= f(x_i) = F'(x) \\ B &= g(x_i) = G'(x) \end{aligned} \tag{5}$$

If you want a number for the complete block, this works:

`\begin{equation}\begin{split}...\end{split}\end{equation}`

$$\begin{aligned} A &= f(x_i) = F'(x) \\ B &= g(x_i) = G'(x) \end{aligned} \tag{6}$$

If you prefer numbering of equations in the form (2.1), (2.2),..., add the line

`\numberwithin{equation}{section}`

to the preamble of your document.

2.2. Theorems and alike. For theorems, lemmas, definitions, etc. use the standard syntax.

`\begin{theorem}...\end{theorem}`

Put optional arguments into square brackets (“Main Theorem” in the example below).

Theorem 2.1 (Main Theorem). *If a knot K has Seifert form V_K and its Alexander polynomial is not 1, then there is an infinite family $\{K_i\}$ of non-concordant knots such that each K_i has Seifert form V_K .*

Definition 2.2. A *preference order* (or *preference relation*) on \mathcal{X} is a binary relation \succ with the following two properties.

1. *Asymmetry:* If $x \succ y$, then $y \not\succeq x$.
2. *Negative transitivity:* If $x \succ y$ and $z \in \mathcal{X}$, then either $x \succ z$ or $z \succ y$ or both must hold.

In this example file, enumerations of theorems, lemmas definitions, etc. appear consecutively. If you want separate numbering (Theorem 2.1, Definition 2.1) change e.g.

```
\newtheorem[theorem]{definition}
```

to

```
\newtheorem{definition}{Definition}[section]
```

If you want a statement unnumbered, just define

```
\newtheorem*{coro}{Corollary}
```

to obtain

Corollary. *If all the coefficients of (A.2) are entire functions, then all local solutions of (A.2) admit a meromorphic continuation over the whole complex plane \mathbb{C} .*

For a proof, use

```
\begin{proof}... \end{proof}
```

An end-of-proof sign \square is set automatically.

Proof. This finishes the proof of the corollary. \square

2.3. Operator names. There are several $\text{T}_{\text{E}}\text{X}$ -commands setting things automatically upright like \det , \sin ,... . If you need operators not predefined, simply define e.g.

```
\newcommand{\Hom}{\operatorname{Hom}}
```

```
\newcommand{\Ker}{\operatorname{Ker}}
```

and then use

```
\Hom, \Ker
```

to obtain

$$\varphi \in \text{Hom}(G/H) \implies \text{Ker}(\varphi) \neq \{0\}.$$

It is desirable that abbreviated mathematical expressions standing for “words” appear in roman (upright) typeface.

3. References

It follows a list of references showing you the style in which authors, books and journal articles should be listed.

- [1] Anderson, M. T., Geometric aspects of the AdS/CFT correspondence. In *AdS/CFT Correspondence: Einstein Metrics and their Conformal Boundaries* (ed. by Olivier Biquard). IRMA Lect. Math. Theor. Phys. 8, European Math. Soc. Publishing House, Zürich 2005, 1–31.
- [2] Babuška, I., Rheinboldt, W. C., Error Estimates for Adaptive Finite Element Computations, *SIAM J. Numer. Anal.* **15** (1978), 736–754.
- [3] Freedman, M. H., Quinn, F., *Topology of 4-manifolds*. Princeton Mathematical Series 39, Princeton University Press, Princeton, NJ, 1990.
- [4] Mal'cev, A. I., On homomorphisms onto finite groups. *Ivanov. Gos. Ped. Inst. Ucen. Zap.* **18** (1958), 49–60.
- [5] Schweizer, M., Hedging of options in a general semimartingale model. Diss. ETH Zürich No. 8615, 1988.
- [7] Vigneras, M. F., Induced representations of reductive p -adic groups in characteristic $\ell \neq p$. *Selecta Math.* (N.S) **4** (1998), 549–623.

Author's address

E-mail: e-mail address